

Choose from the following list 4 exercises that you have not done before, for which you hand in solutions.

Let G be a group with multiplication $m : G \times G \rightarrow G$, inversion $i : G \rightarrow G$ and neutral element e .

Exercise 1 (Isomorphisms, monomorphisms and epimorphisms).

Let $f : G \rightarrow H$ be a group homomorphism. Show that f is an isomorphism (in the sense of Definition 2.3.1) if and only if f is bijective. Show that f is a monomorphism if and only if f is injective. Show that f is an epimorphism if and only if f is surjective.

Exercise 2 (Subgroups).

Let H be a subset of G . Show that H is a subgroup of G if and only if $e \in H$, $m(H \times H) \subset H$ and $i(H) \subset H$. In other words, H is a subgroup if and only if it is a group with respect to the restrictions of m and i to H .

Exercise 3 (The center).

Show that the *center* of G

$$Z(G) = \{ a \in G \mid ab = ba \text{ for all } b \in G \}$$

is a subgroup of G . Show that $Z(G)$ is commutative. Show that every subgroup of $Z(G)$ is normal in G . Is every commutative subgroup of G normal?

Exercise 4 (The subgroup generated by a subset).

1. Let $\{H_i\}_{i \in I}$ be a family of subgroups of G . Show that the intersection $\bigcap_{i \in I} H_i$ is a subgroup of G .
2. Let $S \subset G$ be a subset. Show that

$$\bigcap_{H < G \text{ with } S \subset H} H = \{ a_1 a_2^{-1} \cdots a_{2n-1} a_{2n}^{-1} \mid n \geq 1 \text{ and } a_1, \dots, a_n \in S \cup \{e\} \}$$

and conclude that there is a unique smallest subgroup $\langle S \rangle$ of G that contains S .

Exercise 5 (Orders of elements in commutative groups).

Let G be a commutative group and $a, b \in G$. Show that $\text{ord}(ab)$ divides $\text{ord}(a) \cdot \text{ord}(b)$. Is this also true if G is not commutative?

Exercise 6 (Cyclic groups and the Klein four-group).

1. Classify all cyclic groups up to isomorphism. Which of them are commutative?
2. Show that a cyclic group of order n has a unique subgroup of order d for each divisor d of n .
3. Is the *Klein four-group* $V = (\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z})$ cyclic? Is it commutative?

Exercise 7 (Dihedral groups).

Let D_n be the group of symmetries of a regular polygon with n sides. Show that $D_n = \langle r, s \rangle$ where r is a rotation around the center of the polygon by an angle of $2\pi/n$ and s is the reflection at a line passing through the center of the polygon and one of its vertices. What is the number of elements of D_n ? Show that $D_3 \simeq S_3$, and that for $n \geq 4$, the dihedral group D_n is not isomorphic to a symmetric group.

Exercise 8 (Symmetric groups).

The *symmetric group* S_n is the group of permutations of the numbers $1, \dots, n$, together with composition as multiplication, i.e. $\sigma \cdot \tau = \sigma \circ \tau$. An element σ of S_n is called a *cycle (of length l)* if $\text{ord}(\sigma) = l$ and if there is an $i \in \{1, \dots, n\}$ such that $\sigma(j) = j$ if $j \notin \{i, \sigma(i), \dots, \sigma^{l-1}(i)\}$; we write $\sigma = (i, \sigma(i), \dots, \sigma^{l-1}(i))$ in this case.

1. Show that $(i, \dots, \sigma^{l-1}(i)) = (j, \dots, \sigma^{l-1}(j))$ if $j = \sigma^n(i)$ for some $n \geq 0$.
2. Two cycles $\sigma = (i, \dots, \sigma^{l-1}(i))$ and $\tau = (j, \dots, \tau^{k-1}(j))$ are called *disjoint* if the sets $\{i, \dots, \sigma^{l-1}(i)\}$ and $\{j, \dots, \tau^{k-1}(j)\}$ are disjoint. Show that σ and τ are disjoint if and only if $\sigma\tau = \tau\sigma$.
3. Show that every element of S_n can be written as a product of disjoint cycles.
4. A *transposition* is a cycle (i, j) of length 2. Show that every element of S_n can be written as a product of transpositions.

Exercise 9 (The sign).

Let σ be an element of S_n and $\sigma = \tau_n \circ \dots \circ \tau_1$ and $\sigma = \tau'_m \circ \dots \circ \tau'_1$ two representations of σ as a product of transpositions τ_1, \dots, τ_n and τ'_1, \dots, τ'_m .

1. Show that $n - m$ is even. Conclude that the map $\text{sign} : S_n \rightarrow \{\pm 1\}$ that sends σ to $(-1)^n$ is well-defined.
2. Show that sign is a group homomorphism.

Exercise 10 (Theorem of Cayley).

Let $G = \{a_1, \dots, a_n\}$ be of finite order n . Define the map $f : G \rightarrow S_n$ that sends a_l to the permutation σ_l with $\sigma_l(i) = j$ such that $a_l a_i = a_j$. Show that f is an injective group homomorphism. Conclude that every finite group is isomorphic to a subgroup of a symmetric group.

Exercise 11 (The alternating group).

The *alternating group* A_n is defined as the kernel of $\text{sign} : S_n \rightarrow \{\pm 1\}$. A group G is called *simple* if $G \neq \{e\}$ and if the only normal subgroups of G are $\{e\}$ and G .

1. Show that a cyclic group G of order n is simple if and only if n is a prime number.
2. Show that A_3 is simple. Show that A_4 is not simple. What about A_1 and A_2 ?
3. Show that A_n is simple for $n \geq 5$.¹

Exercise 12 (Quaternion group).

The quaternion group Q consists of the elements $\{\pm 1, \pm i, \pm j, \pm k\}$, and the multiplication is determined by the following rules: 1 is the neutral element, $(-1)^2 = 1$ and

$$i^2 = j^2 = k^2 = -1, \quad (-1)i = -i, \quad (-1)j = -j, \quad (-1)k = -k, \quad ij = k = -ji.$$

1. Is Q commutative?
2. Describe all subgroups of Q .
3. Which subgroups are normal? What are the respective quotient groups?

Exercise 13.

Classify all groups with 6 elements and all groups with 8 elements up to isomorphism.

Exercise 14 (Transitivity of index).

Let H be a subgroup of G and K a subgroup of H . Show that $(G : K) = (G : H)(H : K)$.

Exercise 15 (Quotients by non-normal subgroups).

Let H be subgroup of G . Show that the association $([a], [b]) \mapsto [ab]$ is not well-defined on cosets $[a], [b] \in G/H$ if H is not normal in G .

Exercise 16 (Alternative characterization of normal subgroups).

A subgroup H of G is normal if and only if $gHg^{-1} \subset H$ for every $g \in G$.

Exercise 17 (Exercises on normal subgroups).

Show the following statements.

1. Every subgroup of index 2 is normal.
2. Every subgroup of a commutative group is normal. Is there a non-commutative group G such that every subgroup H of G is normal?
3. The intersection of two normal subgroups is a normal subgroup. If both normal subgroups have finite index, then their intersection has also finite index.

¹This exercise is more difficult than others, but solutions can be found in the literature.

Exercise 18 (Universal property of the quotient).

Let N be a normal subgroup of G . Show that the quotient map $\pi : G \rightarrow G/N$ satisfies the following universal property: for every group homomorphism $f : G \rightarrow H$ with $f(a) = e$ for $a \in N$ there exists a unique group homomorphism $\bar{f} : G/N \rightarrow H$ such that $f = \bar{f} \circ \pi$, i.e. the diagram

$$\begin{array}{ccc} G & \xrightarrow{f} & H \\ \pi \downarrow & \circlearrowleft & \nearrow \bar{f} \\ G/N & & \end{array}$$

commutes.

Exercise 19 (Universal property of the product).

Let $\{G_i\}_{i \in I}$ be a family of groups and $G = \prod G_i$ their product.

1. Show that the map $\pi_i : G \rightarrow G_i$ that sends $(g_i)_{i \in I}$ to g_i is a surjective group homomorphism for every $i \in I$. These maps are called the *canonical projections*.
2. Show that the product together with the canonical projections satisfies the following universal property: for every family of group homomorphisms $\{f_i : H \rightarrow G_i\}_{i \in I}$, there is a unique group homomorphism $f : H \rightarrow \prod G_i$ such that $f_j = \pi_j \circ f$ for every $j \in I$, i.e. the diagram

$$\begin{array}{ccc} H & \xrightarrow{f} & \prod G_i \\ & \searrow f_i & \downarrow \pi_j \\ & & G_j \end{array}$$

commutes for every $j \in I$.

Exercise 20 (Universal property of the direct sum).

Let $\{G_i\}_{i \in I}$ be a family of commutative groups and $G = \bigoplus G_i$ their direct sum.

1. Show that the map $\iota_i : G_i \rightarrow G$ that sends g to $(g_j)_{j \in I}$ with $g_i = g$ and $g_j = e_j$ for $j \neq i$ is an injective group homomorphism for every $i \in I$. These maps are called the *canonical injections*.
2. Show that the direct sum together with the canonical injections satisfies the following universal property: for every family of group homomorphisms $\{f_i : G_i \rightarrow H\}_{i \in I}$ of commutative groups, there is a unique group homomorphism $f : \bigoplus G_i \rightarrow H$ such that $f_j = f \circ \iota_j$ for every $j \in I$, i.e. the diagram

$$\begin{array}{ccc} \bigoplus G_i & \xrightarrow{f} & H \\ \iota_j \uparrow & \circlearrowleft & \nearrow f_j \\ G_j & & \end{array}$$

commutes for every $j \in I$.

3. Is the same true if H is a non-commutative group?

Exercise 21 (Some group actions).

Show that the following maps are group actions.

1. $S_n \times \{1, \dots, n\} \rightarrow \{1, \dots, n\}$, with $\sigma.i = \sigma(i)$.
2. $\text{GL}_n(\mathbb{R}) \times \mathbb{R}^n \rightarrow \mathbb{R}^n$, with $g.v = g \cdot v$ (usual matrix multiplication).
3. $\mathbb{R}^\times \times \mathbb{R}^n \rightarrow \mathbb{R}^n$, with $a.v = a \cdot v$ (scalar multiplication).
4. The permutation of the vertices of a regular n -gon by elements of the dihedral group D_n .

Exercise 22 (Center and centralizer).

Consider the action of G on itself by conjugation.

1. Show that

$$\{x \in G \mid \mathcal{O}(x) = \{x\}\} = \{a \in G \mid ab = ba \text{ for all } b \in G\}.$$

2. Show that $C_G(x) = \{a \in G \mid ax = xa\}$.
3. Show that

$$Z(G) = \bigcap_{x \in G} C_G(x).$$

Exercise 23 (Normalizer).

Let H be a subgroup of G . Show that its normalizer $\text{Norm}_G(H)$ is the largest subgroup of G containing H such that H is a normal subgroup of $\text{Norm}_G(H)$. Show further that the following properties are equivalent:

1. H is normal in G ;
2. $\text{Norm}_G(H) = G$;
3. H is a fixed point for the action of G on the set of all subgroups of G by conjugation.

Exercise 24 (Short exact sequences).

A short exact sequence of groups is a sequence

$$\{e\} \xrightarrow{f_1} N \xrightarrow{f_2} G \xrightarrow{f_3} Q \xrightarrow{f_4} \{e\}$$

of groups and group homomorphism such that $\text{im} f_i = \ker f_{i+1}$ for $i = 1, 2, 3$.

1. Show that $\text{im} f_i = \ker f_{i+1}$ for $i = 1, 2, 3$ holds if and only if f_2 is injective, if $\text{im} f_2 = \ker f_3$ and if f_3 is surjective.
2. Show that N is isomorphic to $N' = \text{im} f_1$, that N' is a normal subgroup of G and that $G/N' \simeq Q$ in case of a short exact sequence.

Exercise 25.

Calculate all orbits and stabilizers for the action of D_4 on itself by conjugation.

Exercise 26 (Commutator subgroup).

The commutator of two elements $a, b \in G$ is $[a, b] = aba^{-1}b^{-1}$. The commutator subgroup of G is the subgroup $[G, G]$ generated by the commutators $[a, b]$ of all pairs of elements a and b of G .

1. Show that $[a, b] = e$ if and only if $ab = ba$. Conclude that $[G, G] = \{e\}$ if and only if G is commutative.
2. Show that $c[a, b]c^{-1} = [cac^{-1}, cbc^{-1}]$ and conclude that $[G, G]$ is a normal subgroup of G .
3. Show that the quotient group $G^{\text{ab}} = G/[G, G]$ is commutative.
4. Show that G^{ab} together with the projection $\pi : G \rightarrow G^{\text{ab}}$ satisfies the following universal property: for every group homomorphism $f : G \rightarrow H$ into a commutative group H , there exists a unique group homomorphism $f^{\text{ab}} : G^{\text{ab}} \rightarrow H$ such that $f = f^{\text{ab}} \circ \pi$:

$$\begin{array}{ccc}
 G & \xrightarrow{f} & H \\
 \pi \downarrow & \circlearrowleft & \nearrow f^{\text{ab}} \\
 G^{\text{ab}} & &
 \end{array}$$

Exercise 27.

Determine all p -Sylow subgroups of S_4 for $p \in \{2, 3\}$.

Exercise 28.

Let $\text{ord}(G) = 6$ and n_p the number of p -Sylow subgroups of G . Find all possibilities for n_2 and n_3 , using the Sylow theorems. Find examples of groups with 6 elements that realize these possibilities.

Exercise 29.

Let $\text{ord}(G) = pq$ for prime numbers p and q . Show that G is not simple.

Hint: If $p = q$, then use the class equation. If $p \neq q$, then use the Sylow theorems.